



Dental Clinical Policy

Subject: Osseous Surgery (Periodontal)

Guideline #: 04-205

Status: Revised

Publish Date: 01/01/2023

Last Review Date: 10/26/2022

Description

This document addresses the procedure of osseous surgery used in the treatment of periodontal disease when there is resultant bone loss.

The plan performs review of osseous surgery due to contractual requirements that necessitate benefits for dental services meet specific contract requirements. For example, plan contract(s) may require the provision of benefits for services that meet generally accepted standards of dental care at the lowest cost that properly addresses the patient's condition. The conclusion that a particular service is medically or dentally necessary and/or appropriate does not constitute an indication and/or warranty that the service requested is a covered benefit payable by the dental plan.

Clinical Indications

When gingivitis progresses to periodontal disease, osseous surgery may be necessary. For osseous surgery to be appropriate, bone loss (horizontal or vertical defects) must be documented around teeth and dental implants and with pocket depth recordings measuring 5 millimeters or greater with bleeding upon probing.

Dental review as it applies to accepted standards of care means dental services that a Dentist, exercising prudent clinical judgment, provides to a patient for the purpose of evaluating, diagnosing or treating a dental injury or disease or its symptoms, and that are: in accordance with the generally accepted standards of dental practice; in terms of type, frequency and extent and is considered effective for the patient's dental injury or disease; and is not primarily performed for the convenience of the patient or dentist, is not cosmetic and is not more costly than an alternative service.

For dental purposes, "generally accepted standards of dental practice" means:

- Standards that are based on credible scientific evidence published in peer-reviewed, dental literature generally recognized by the practicing dental community
- specialty society recommendations/criteria
- the views of recognized dentists practicing in the relevant clinical area
- any other relevant factors from credible sources

Criteria

Osseous surgery procedures are considered appropriate with:

1. Completion of initial periodontal therapy (e.g. scaling and root planing) allowing a minimum of four weeks prior to any surgical treatment for the tissues to properly heal which allows for proper assessment of the success or failure of non-surgical therapy.
2. Post initial therapy periodontal charting demonstrating pocket depths greater than or equal to 5mm.
3. Current (within 12 months), dated, post initial therapy periodontal charting (6-point periodontal charting) indicating pocket depth recordings of a minimum of 5mm.
4. Current (within 12 months), dated, diagnostic, pretreatment radiographic images demonstrating either horizontal and/or vertical osseous defects.
5. When radiographic images are not demonstrative, a detailed narrative describing periodontal case type, percentage of bone loss, measured furcation involvement, and description of the vertical defect may be requested.
6. Benefits will be limited to two quadrants per date of service. Exceptions will be allowed on a case-by-case basis.
7. Benefits are group contract dependent but are limited to one (1) osseous surgical procedure in a given period per single tooth or multiple teeth in the same quadrant and only if the pocket depth of the tooth is a minimum of 5mm.
8. Periodontal surgical procedures such as, but are not limited to, gingivectomy or gingivoplasty, mesial/distal wedge procedure, anatomical crown exposure, gingival flap procedure, apically repositioned flap, clinical crown lengthening, and surgical revision procedure are considered inclusive with osseous surgery.
9. Only those teeth with favorable periodontal outcomes will be considered for benefit.

Osseous surgery may be appropriate for the treatment of periodontal disease defects on natural teeth and dental implants. Dental implants will be considered the same as a natural tooth as it relates to quadrant surgery.

Osseous surgery post-operative management as well as for any surgical re-entry has a frequency limitation as per group contract.

Note: Whether a service is covered by the plan, when any service is performed in conjunction with or in preparation for a non-covered or denied service, all related services are also either non-covered or denied.

Coding

The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

CDT

Including, but not limited to, the following:

D4260

Osseous surgery (including elevation of a full thickness flap and closure) – four or more contiguous teeth or tooth bounded spaces per quadrant

D4261	Osseous surgery (including elevation of a full thickness flap and closure) – one to three contiguous teeth or tooth bounded spaces per quadrant
D6101	Debridement and osseous recontouring of peri-implant defect or defects surrounding a single implant, surface cleaning of the exposed implant surfaces, including flap entry and closure.
D6102	Debridement and osseous recontouring of a peri-implant defect or defects surrounding a single implant, and includes surface cleaning of the exposed implant surfaces, including flap entry and closure

ICD-10 CM Diagnoses for Dental Diseases and Conditions: See the current CDT code book for details

References:

1. Proestakis G, Soderholm G, et al. Gingivectomy versus flap surgery: the effect of the treatment of infra-bony defects. A clinical and radiographic study. *Journal of Clinical Perio* 1992a; 19:497-508.
2. Bower RC. Furcation morphology relative to periodontal treatment furcation entrance architecture. *J Perio* 1979; 50:23-27.
3. Waerhaug J. Healing of the dento-epithelial junction following sub-gingival plaque control. *Journal of Perio* 1978; 49:119-134.
4. Proceedings of the World Workshop in clinical Periodontics. Sec II Non- surgical Periodontal Treatment. II-4 and II-15. July, 1989 American Academy of Periodontology. 211 E Chicago Ave., Chicago, IL.
5. Parameter on Chronic Periodontitis with Advanced Loss of Periodontal Support. Parameters of Care. Supplement *J Perio* 2000; 71: 856-858.
6. Rabinni GM, Ash MA and Caffesse RG. The effectiveness of scaling and root planing in calculus removal. *J Perio* 1981; 52:119.
7. Parashis AO, Anagnou-Vareltzides A and Demetriou, N. Calculus Removal from multi-rooted teeth with and without surgical access. *Journal of Clinical Perio* 1993; 20: 63-68.
8. Prichard JP. The infrabony technique as a predictable procedure. *Journal of Perio* 1957; 28:202.
9. Parameter on Chronic Periodontitis with Slight to Moderate Loss of Periodontal Support Parameters of Care. Supplement *J Perio* 2000, 71: 853-855.
10. Persson R and Swindsen J. The role of periodontal probing depth in clinical decision making. *Journal of Clinical Perio* 1990; 17:96-101.
11. CDT 2023 Current Dental Terminology, American Dental Association
12. Treatment of Plaque Induced Gingivitis, Chronic Periodontitis, and Other Clinical Conditions. Academy Report. American Academy of Periodontology. *J Perio* 2001, 72: 1790-1800.
13. Palcanis KG. Surgical pocket therapy. *Annals of Periodontology* 1996; 1:589-617.
14. Greenstein G. Non-surgical periodontal therapy in 2000: a literature review. *JADA* 2000; 131:1580-1592.
15. American Academy of Periodontology. Treatment of gingivitis and periodontitis (position paper). *Amer Acad Perio*. Dec 1997.
16. Buchanan SA and Robertson PA. Calculus removal by scaling/root planing with and without surgical access. *J Perio* 1987; 58:159-163.
17. Caffesse RG, Sweeney PL and Smith BA. Scaling and root planing with and without periodontal flap surgery. *J Clin Perio* 1986; 13:205-210.

18. Journal of the American Dental Association. Consensus reports from the 1996 world workshop in periodontics. JADA supplement 1996; 129:40-S, 41-s.
19. Gher ME and Vernino AR. Root anatomy: a factor in inflammatory periodontal disease. Int'l J Perio and Restorative Dentistry 1981; 1(5):63-68.
20. Claffey N. Decision making in periodontal therapy: The reevaluation. J Clin Periodontal 18:364, 1991.

History

Revision History	Version	Date	Nature of Change	SME
	initial	12/14/16	creation	M Kahn G Koumaras
	Revision	2/8/17	General verbiage	Rosen
	Revision	2/16/18	Appropriateness/medical necessity, criteria	M Kahn
	Revision	10/08/2020	Annual Review	Committee
	Revised	12/4/2020	Annual Review	Committee
	Revised	10/30/2021	Annual Review	Committee
	Revised	10/26/2022	Annual Review	Committee

Federal and State law, as well as contract language, takes precedence over Dental Clinical Policy. Dental Clinical Policy provides guidance in interpreting dental benefit application. The Plan reserves the right to modify its Dental Clinical Polices and guidelines periodically and as necessary. Dental Clinical Policy is provided for informational purposes and does not constitute medical advice. These Policies are available for general adoption by any lines of business for consistent review of the medical or dental necessity and/or appropriateness of care of dental services. To determine if a review is required, please contact the customer service number on the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the Plan.

Current Dental Terminology - CDT © 2023 American Dental Association. All rights reserved.